



ONION

2017 RANGE BROCHURE

Ph: 1300 LEFROY (533 769)

www.lefroyvalley.com

Onion (*Allium cepa* L.)

Variety	Day Length	Colour	Storage Potential	Sowing
Brown				
CARBIDE	Intermediate	Bronze brown	6-7 months	May to early June
KODAF^{F1}	Short	Bronze brown	3-4 months	Mid May to end of May in QLD, Last week in May to June in NSW and South Australia.
LEGION^{F1}	Short	Bronze brown	2-3 months	Mid May to end of May in QLD, May to June in NSW and SA
SALUTE^{F1}	Intermediate	Golden copper	3-4 months	Early to main season
SPARTAN^{F1}	Short	Bronze brown	2-3 months	Mid march to April
White				
CUMULUS^{F1}	Intermediate	White	3 to 4 months	July to September
NIMBUS^{F1}	Long	White	4 to 5 months	July to September
Red				
RUBY ORB^{F1}	Long	Red	4 to 5 months	Mid August to end of September



Cumulus^{F1}



Nimbus^{F1}

Onion Bunching

Variety	Plant	Colour	Harvest Season
ARCTIC	Erect uniform plant	Medium dark green	Colour is darker during summer. Can be used in all production windows IR: Pd
ICICLE^{F1}	Erect uniform plant	Dark green	Can be used in all production windows. Best suited to winter production IR to Pd



Arctic^{F1}



Icicle^{F1}

DISEASE KEY DEFINITIONS

HR: Highly resistant plant varieties would highly restrict the growth and development of the specified pest or pathogen under normal growing conditions and pest pressure when compared to susceptible varieties. These plant varieties may, however, exhibit some symptoms or damage under heavy pest pressure or stressful growing conditions.

IR: Intermediately resistant plant varieties would restrict the growth and development of the specified pest or pathogen. The plant variety may however exhibit a greater range or degree of symptoms when compared to a highly resistant variety. Intermediately resistant plant varieties will still show less severe symptoms or disease damage than susceptible plant varieties when the crop is grown under similar environmental conditions and is subjected to the same disease or pest pressure.

Pd - Downy mildew (*Peronospora destructor*)

DISCLAIMER

Lefroy Valley carries out trialling throughout Australia/New Zealand prior to releasing varieties into the market place. We strongly recommend that all varieties be trialed under your growing conditions prior to commercial sowings taking place. For details of up-to-date technical information & trial results in your area please contact Lefroy Valley. This information is valid at the date of publication. All cultural and descriptive information is supplied in good faith as a guide only. Varietal performance is influenced by many variables, namely climatic, soil conditions, cultural and management practices. No liability will be accepted by Lefroy Valley or its representatives as to final performance based on this information.



1300 LEFROY (533 769)

VIC/SA/TAS

PH: 03 8779 2121
FX: 03 8732 0308

WA/NT

PH: 08 9344 4000
FX: 08 6210 1360

QLD/NSW

PH: 07 4632 0555
FX: 07 4632 0155

orders@lefroyvalley.com

www.lefroyvalley.com

Publication Date: December 2016